

A New Sufficient Condition for Uniqueness in Continuous Games

Seamus Hogan*
Department of Economics
University of Canterbury
seamus.hogan@canterbury.ac.nz

Abstract:

Consider the class of games in which each player chooses a strategy from a connected subset of the real line. Many oligopoly models fall into this class. In many of these applications, it would be useful to show that an equilibrium was unique, or at least to have a set of conditions under which uniqueness would hold.

In this paper, we first prove a uniqueness theorem that is slightly less restrictive than the contraction mapping theorem for mappings from the subsets of the real line onto itself, and then show how uniqueness in the general game can be shown by proving uniqueness using an iterative sequence of P to P mappings. This iterative approach works by considering the equilibrium for an m -player game holding the strategies of all other players fixed, starting with a two-player game. If one can show that the m -player game has a unique equilibrium for all possible values for the remaining players strategies, then one can add one player at a time and consider the P to P mapping from that player's strategy on to the unique equilibrium of the first m players and back onto the $(m+1)$ th palyer's strategy.

We then show how a general condition for each one of this sequence of mappings to have a unique equilibrium is that the Jacobean matrix of best-response functions be positive semi definite, and how this general condition encompasses and generalizes some existing uniqueness theorems for particular games.

* I would like to thank Douglas Bridges, Clive Granger, Ngo van Long, Jan Magnus, Les Oxley, and Asha Saddanand for helpful comments on earlier versions of this paper.

A New Sufficient Condition for Uniqueness in Continuous Games

1. Introduction:

Many oligopoly models fall into the class of continuous games in which each player chooses a strategy from a connected subset of the real line. For example, in Cournot models, each firm chooses a quantity to produce, in product differentiation models firms typically choose a price, and so on. Existence of a Nash equilibrium in pure strategies in such models are generally shown by means of the Brouwer or Kakutani fixed-point theorems. In many of these applications, it would be useful to show that an equilibrium was unique, or at least to have a set of easy-to-apply conditions under which uniqueness would hold.

There are a number of uniqueness conditions that have been found for particular subsets of this general class of models.¹ Generally, these involve a trade-off between generality and ease of application. For instance, the classic paper by Rosen (1965) offers a general sufficient condition for uniqueness in a wide class of games that encompasses the class considered in this paper, but the condition is quite opaque and hence difficult to apply directly to particular applications. At the opposite end of the generality-usability continuum, are uniqueness conditions that are specific to particular applications. For instance, conditions for a unique equilibrium in the Cournot quantity-setting oligopoly model have been derived by Szidarovszky and Yakowitz (1977), Gaudet and Salant (1991), and Long and Soubeyran (2000).

One of the most useful general ways of showing uniqueness is through the contraction mapping theorem. The contraction mapping theorem provides a quite general condition for there to be a unique fixed point, with the added benefit that it guarantees existence without the requirements that the space being mapped onto itself be convex or bounded. To be useable, however, one needs to show that the particular mapping for which a fixed point defines an equilibrium constitutes a contraction mapping, a task that is not always straightforward.

In this paper, we derive a variant of the contraction mapping theorem, and present an approach in which this variant is applied iteratively to the class of games outlined above to derive an easy-to-apply uniqueness condition defined in terms of the slopes of the players' best-response functions. This uniqueness condition encompasses and generalises a number of existing uniqueness conditions.

In the next section, we present the contraction mapping theorem and a related, less-restrictive theorem for the special case of a mapping from a subset of the real line onto itself. Section 3 lays out the general problem and shows by example why the conventional characterisation of an equilibrium as a fixed point of an \mathbb{R}^n -to- \mathbb{R}^n mapping is too restrictive. Section 4 shows how an equilibrium in the general game can be defined in terms of a sequence of contraction mappings involving \mathbb{R} -to- \mathbb{R} mappings; Section 5 then shows how the uniqueness condition derived iteratively in this way can be represented in terms of the slopes of the best-response functions of each player. Finally, Section 6 shows how this general uniqueness condition encompasses and generalises many existing results.

¹ A good survey of the existing uniqueness theorems is contained in Cachon and Netessine (2004).

2. The Contraction Mapping Theorem and a Related Result.

A. The Contraction Mapping Theorem in Euclidean Space.

Typically in oligopoly models, the existence of an equilibrium is proved by showing the existence of a fixed point in a mapping from a subset of Euclidean space onto itself. Let X be a subset of \mathbb{R}^n , and let $f : X \rightarrow X$ be a single-valued function mapping X onto itself. In this context, the definition of a contraction mapping and the contraction mapping theorem are as follows:

Definition 1:

If there exists $b \in (0,1)$ and a norm $\|\cdot\|$ such that

$$\|f(y) - f(x)\| \leq b \|y - x\| \quad \forall x, y \in X, \quad (1)$$

then f is a contraction mapping.

Theorem 1 (The Contraction Mapping Theorem):

If X is a closed subset of \mathbb{R}^n and f is a contraction mapping, then

- a) (existence and uniqueness) there exists a unique fixed point $x^* \in X$ such that $f(x^*) = x^*$,
- b) (convergence) for any $x \in X$ and $n \geq 1$, $\|f^n(x) - x^*\| \leq b^n \|x - x^*\|$.

The contraction mapping theorem has three advantages over the Brouwer or Kakutani fixed point-theorems if a contraction mapping can be shown to exist: First, and most importantly, it shows uniqueness as well as existence; second, it does not require that the set X be bounded; and third, it has the convergence property.

The convergence property implies that the unique fixed point can easily be found numerically. It can be useful in a game-theoretic context if we imagine the Nash equilibrium to be one iteration of a repeated game, as it suggests that the Nash-equilibrium outcome can be stable in the sense that if every period each player chooses the best response to the previous-period strategies of the other players, the game will converge to the unique equilibrium.

Such dynamic interpretations of a static equilibrium are not always appropriate, however, and numerical solveability is not often important. If we only require existence and uniqueness and not convergence, we can, in principle, relax Condition (1). We do this below for the case of \mathbb{R} -to- \mathbb{R} mappings.

B. The Contraction Mapping Theorem in \mathbb{R}^1 Space.

In this paper, we show how existence of an equilibrium that is a point in Euclidean n space, can be represented as a set of fixed points of a sequence of mappings from the real line onto itself. For \mathbb{R} -to- \mathbb{R} mappings, the natural norm to use is the absolute value, $\|x\| = |x|$, and the definition of a contraction mapping becomes as follows:

Definition 2:

If there exists $b \in (0,1)$ such that

$$\frac{|f(y) - f(x)|}{|y - x|} \leq b \quad \forall x, y \in X, \quad (2)$$

then f is a contraction mapping.

In words, this says that the straight line between any two points on the graph of the function, must have a slope in the interval $(-1,1)$. If, we don't require the convergence property, we only require that the slope be less than 1. We will define such a function as a "quasi-contraction mapping".

Definition 3:

If there exists $b \in (0,1)$ such that

$$\frac{f(x) - f(y)}{x - y} \leq b \quad \forall x, y \in X, \quad (3)$$

then f is a quasi-contraction mapping.

This gives us the following variant of the contraction mapping theorem:

Theorem 2:

If X is a closed subset of \mathbb{R} and f is a quasi-contraction mapping, then there exists a unique fixed point, $x^* \in X$, such that $f(x^*) = x^*$.

Proof:

First we show that a fixed point must exist. For any $x_0 \in X$, we have $f(x_0) \geq x_0$, or $f(x_0) \leq x_0$. If $f(x_0) \geq x_0$, then define,

$$y_0 = \begin{cases} \max\{x | x \in X\} & \text{if } X \text{ is bounded above} \\ \frac{f(x_0) - b x_0}{1 - b} & \text{otherwise} \end{cases} \quad (4)$$

If X is not bounded above, we have, from (3) and (4),

$$\begin{aligned} f(y_0) - f(x_0) &\leq b(y_0 - x_0) \\ \Rightarrow f(y_0) &\leq b y_0 + f(x_0) - b x_0 \\ \Rightarrow f(y_0) &\leq y_0. \end{aligned}$$

If X is not bounded above, we have directly that

$$f(y_0) \leq y_0.$$

By the intermediate value theorem, therefore, there exists $x^* \in [x_0, y_0]$ such that $f(x^*) = x^*$. A similar argument holds if $f(x_0) \leq x_0$.

To show uniqueness, let $x^* \in X$ be a fixed point of f . Then, $\forall x > x^*$,

$$\frac{f(x) - f(x^*)}{x - x^*} \leq b < 1$$

$$\Rightarrow f(x) - f(x^*) \leq b(x - x^*) < x - x^*$$

$$\Rightarrow f(x) - x < f(x^*) - x^* = 0.$$

Thus any fixed point of f must be the maximum fixed point, implying that only one can exist.

Ω

Finally, if, in addition to the above assumptions, we assume that X is a connected subset of \mathbb{R} and that f is differentiable almost everywhere, then Condition (2) is equivalent to the following:

- a) f is continuous over X ,
 - b) $|f'(x)| \leq b \quad \forall x \in X$ where f is differentiable.
- (5)

Similarly, Condition (3) is equivalent to the following:

- a) f is continuous over X ,
 - b) $f'(x) \leq b \quad \forall x \in X$ where f is differentiable.
- (6)

This gives the general uniqueness theorem we shall use in this paper:

Theorem 3:

Let X be a closed, connected subset of \mathbb{R} , and let f be a single-valued continuous function from X onto itself that is differentiable almost everywhere. Then if, for some $b \in (0, 1)$

- a) $f'(x) < b \quad \forall x \in X$ where f is differentiable,

there exists a unique fixed point $x^* \in X$ such that $f(x^*) = x^*$.

If in addition, we have

- b) $f'(x) > -b \quad \forall x \in X$ where f is differentiable,

then the fixed point is stable in the sense that for any $x \in X$ and $n \geq 1$, $|f^n(x) - x^*| \leq b^n |x - x^*|$.

In the next section we show how the contraction mapping theorem in \mathbb{R}^n space is used to establish uniqueness in the class of games considered in this paper, and show by example why we seek to reduce the problem to one involving \mathbb{R} -to- \mathbb{R} mappings.

3. The General Problem:

A. Notation:

Imagine that there are n players. We employ the following notation. The strategy space for each player i is X_i , and the set of all possible combinations of strategies for all players is

X^n . Let $x_i \in X_i$ denote a strategy for player i and let $\mathbf{x} = (x_1, \dots, x_n) \in X^n$ denote a strategy combination for all players. Rather than specify the payoffs for each player, we will express everything in terms of the best-response functions. Let \mathbf{x}_{-i} denote the $(n-1)$ -vector of strategies of all players other than player i , and let $f_i(\mathbf{x}_{-i})$ be the best-response of player i to this combination of strategies of the other players.

We impose the following restrictions on this general set-up:

Assumption 1:

- a) For each i , X_i is a connected subset of the real line;
- b) for each i , f_i is continuous, single-valued, and differentiable almost everywhere over X^n .

We do not require that the f_i be fully differentiable so that the model will be able to handle non-differentiabilities that can arise from boundary solutions to an individual player's optimisation problem. For ease of exposition, however, when presenting expressions involving derivatives we will omit the repeated caveat, " $\forall \mathbf{x} \in X^n$ where \mathbf{f} is differentiable", but this is implied.

Proofs of existence of an equilibrium in this class of games typically proceed by defining the aggregate best-response function, $\mathbf{f} : X \rightarrow X$ where $\mathbf{f} = (f_1, f_2, \dots, f_n)$, so that a Nash equilibrium in pure strategies is a fixed point of \mathbf{f} and vice versa, and then appealing to the Brouwer fixed-point theorem. We want to find sufficient conditions for \mathbf{f} to have a unique fixed point. For this, it would be sufficient to show that \mathbf{f} is a contraction mapping. This, however, would be too restrictive. This is illustrated by the following simple example.

B. A Numerical Example:

Consider a Cournot game in which the n players are firms choosing the quantity to produced taking the quantity produced by each of the other $n-1$ firms as given. Assume that the market inverse demand curve is linear, and that each firm has a constant marginal cost of production. The linear structure satisfies the conditions required for uniqueness by Szidarovszky and Yakowitz (1977) amongst others.

To see whether the aggregate best-response function constitutes a contraction mapping, we need to define a norm. Rather than choosing a particular norm, we will consider any p -norm of the form

$$\|\mathbf{x}\|_p = \left(\sum_{i=1}^n (|x_i|)^p \right)^{\frac{1}{p}} \text{ for some real number } p \geq 1.$$

Now consider some initial vector of outputs, $\mathbf{x} = (x_1, \dots, x_n)$, and a second vector, $\mathbf{y} = (y_1, \dots, y_n)$, where all outputs have been perturbed by the same constant, d . $y_i = x_i + d \quad \forall i$. The p -norm for this perturbation is

$$\|\mathbf{y} - \mathbf{x}\|_p = n^{1/p} d.$$

The linear Cournot game produces linear best-response functions in which, at interior solutions, $\partial x_i / \partial x_j = -0.5 \quad \forall i, j \neq i$. We therefore have $f_i(\mathbf{x}_{-i}) = x_i - 0.5(n-1)d$, and hence

$$\|\mathbf{f}(\mathbf{y}) - \mathbf{f}(\mathbf{x})\|_p = n^{1/p} 0.5(n-1)d.$$

For any $n > 2$, therefore, we have

$$\|\mathbf{f}(\mathbf{y}) - \mathbf{f}(\mathbf{x})\|_p \geq \|\mathbf{y} - \mathbf{x}\|_p$$

and hence \mathbf{f} is not a contraction mapping for any p -norm.

What this example shows is that if we wish to use the contraction mapping theorem to show uniqueness in a class of models that encompasses this standard example, we will need to use a different function than \mathbf{f} for which a fixed point defines an equilibrium. In the next section, we develop an approach that enables us to transform the problem so that an equilibrium is a fixed point in a mapping from a subset of the real line onto itself.

4. An Alternative Approach:

To transform the problem, we define an equilibrium iteratively, starting with 2 players and then progressively adding more in. To do this, define an m -equilibrium as a set of x_i such that

$$x_i = f_i(\mathbf{x}_{-i}) \quad \forall i = 1..m,$$

where $m \leq n$. That is, it is a set of strategies such that the first m players' strategies are the best response to the strategies of *all* other players, but the remaining $n-m$ players' strategies are unconstrained.

Let \mathbf{x}^m be the vector of strategies by the first m firms, and let \mathbf{x}^{-m} be the strategies of the remaining $n-m$ firms when $m < n$. An m -equilibrium when $m < n$ is therefore a Nash equilibrium in \mathbf{x}^m taking \mathbf{x}^{-m} as given, and the m -equilibrium when $m = n$ is a Nash equilibrium of the full game. Let $h_i^m(\mathbf{x}^{-m})$ denote an m -equilibrium value of x_i when $m < n$ and let $\mathbf{h}^m(\mathbf{x}^{-m})$ be an m -vector of those values. If there exists a unique m -equilibrium for each value of \mathbf{x}^{-m} , then the function \mathbf{h}^m is single valued and defined for all \mathbf{x}^{-m} .

Our approach to finding sufficient conditions for a unique equilibrium is to find the conditions for a unique 2-equilibrium that holds for all values of \mathbf{x}^{-2} and then to extend that by induction by finding conditions for there to exist a unique $(m+1)$ -equilibrium that holds for all values of $\mathbf{x}^{-(m+1)}$ conditional on there being a unique m -equilibrium.

Let $m=2$. Define the mapping $g_2 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ as

$$g_2(x_2, \mathbf{x}^{-2}) \equiv f_2(f_1(x_2, \mathbf{x}^{-2}), \mathbf{x}^{-2}). \quad (7)$$

There is an equivalence between a 2-equilibrium and a fixed point of g_2 . From Theorem 2, a sufficient condition for there to exist a unique 2-equilibrium is that g_2 be a quasi-contraction mapping.

Now imagine that there exists a unique $(m-1)$ -equilibrium for each value of $\mathbf{x}^{-(m-1)}$. In this case, define the mapping $g_m : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ as

$$g_m(x_m, \mathbf{x}^{-m}) \equiv f_m(\mathbf{h}^{m-1}(\mathbf{x}^{-(m-1)}), \mathbf{x}^{-(m-1)}). \quad (8)$$

Again, there is an equivalence between an m -equilibrium and a fixed point of g_m , and so a sufficient condition for there to exist a unique m -equilibrium given x^{-m} is that g_m be a quasi-contraction mapping.

Definition 4:

We say that f exhibits an “iterative quasi-contraction mapping” if g_m exists and is a quasi-contraction mapping for each $m \in \{2 \dots n\}$, and that it exhibits an “iterative contraction mapping” if g_m exists and is a contraction mapping for each $m \in \{2 \dots n\}$.

The main result of this paper is then

Theorem 4:

If the best-response functions f exhibit an iterative quasi-contraction mapping, then the game has a unique Nash equilibrium.

Proof:

As we have shown, if g_2 is a quasi-contraction mapping, there exists a unique 2-equilibrium for all values of x^{-2} . If there exists a unique $(m-1)$ -equilibrium for all values of $x^{-(m-1)}$ then g_m exists, and if g_m is a quasi-contraction mapping, there exists a unique m -equilibrium. By induction, then, if g_m is a quasi-contraction mapping for each $m \in \{2 \dots n\}$, then there must exist a unique m -equilibrium for each m , and hence a unique equilibrium for the full game.

Ω

Now imagine that each of the g_m is a full contraction mapping so that repeated applications of g_m will generate convergence to the unique fixed point. This does not imply that the full equilibrium would be stable in the way it would be if f were a contraction mapping.² It does, however, that imply a numerical solvability using the best-response functions in the following sense. First, note that the 2-equilibrium can found iteratively by alternately adjusting player 1’s strategy to that of player 2 and vice versa. Then, if the m -equilibrium is iteratively solvable by sequentially adjusting each of the first m player’s strategies to be on his best-response functions, and if g_{m+1} is a contraction mapping, then the $(m+1)$ -equilibrium is iteratively solvable by adjusting the m -equilibrium to x_{m+1} and then x_{m+1} to the h^m and so on. Iterative solveability is perhaps not the most useful property one might desire of an equilibrium, but it is essentially a free result.

5. Sufficient Conditions with Calculus.

The analysis of the previous section gives sufficient conditions for uniqueness and iterative solvability that derive from our sequential approach. They are not, however,

² This can be seen from the example in Section 3, for which the equilibrium is not stable, but for which, as will be shown in Section 6, each of the g_m is a contraction mapping.

particularly user friendly. For that, we would like to express the conditions in terms of the slopes of the best-response functions.

To do this let \mathbf{J} be the $n \times n$ Jacobean matrix of \mathbf{f} with elements J_{ij} , so that

$$J_{ij} = \frac{\partial f_i}{\partial x_j}, \quad \forall j \neq i, \forall i, \quad \text{and} \quad J_{ii} = 0 \quad \forall i.$$

That is, \mathbf{J} is the matrix of slopes of the best-response functions of each player with respect to the strategies of each other player.

Let \mathbf{A}_n be the $n \times n$ matrix, $\mathbf{A}_n = \mathbf{I}_n - \mathbf{J}$, where \mathbf{I}_n is the $n \times n$ identity matrix. Finally, let \mathbf{J}_m and \mathbf{A}_m be the $m \times m$ matrices comprising the first m rows and first m columns of \mathbf{J} and \mathbf{A}_n , respectively (that is, the principal minors of order m of \mathbf{J} and \mathbf{A}_n). The derivatives of the functions g_m can be expressed in terms of the determinants of the \mathbf{A}_m as follows:

Theorem 5:

If g_m exists, then

$$\frac{dg_m}{dx_m} = 1 - \frac{|\mathbf{A}_m|}{|\mathbf{A}_{m-1}|}. \quad (9)$$

Proof:

Note that

$$|\mathbf{A}_1| = 1, \quad \text{and} \quad |\mathbf{A}_2| = 1 - \frac{\partial f_2}{\partial x_1} \cdot \frac{\partial f_1}{\partial x_2},$$

and, from Equation (7),

$$\frac{dg_2}{dx_2} = \frac{\partial f_2}{\partial x_1} \cdot \frac{\partial f_1}{\partial x_2},$$

so Equation (9) holds for $m=2$.

For $m > 2$, we have from Equation (8) that

$$\frac{dg_m}{dx_m} = \sum_{i=1}^{m-1} \frac{\partial f_m}{\partial x_i} \cdot \frac{\partial h_i^{m-1}}{\partial x_m}. \quad (10)$$

Define \mathbf{A}_{Cm} as the column vector containing the first $m-1$ elements of the m 'th column of \mathbf{A} , and define \mathbf{A}_{Rm} similarly as the row vector containing the first $m-1$ elements of the m 'th row of \mathbf{A} , so that \mathbf{A}_m is the partitioned matrix

$$\mathbf{A}_m = \begin{bmatrix} \mathbf{A}_{m-1} & \mathbf{A}_{Cm} \\ \mathbf{A}_{Rm} & 1 \end{bmatrix}.$$

We can then rewrite Equation (10) as

$$\frac{dg_m}{dx_m} = -\mathbf{A}_{Rm} \frac{\partial \mathbf{h}^{m-1}}{\partial x_m}. \quad (11)$$

The h_i^{m-1} variables are defined by the fixed point in the m -equilibrium

$$h_i \equiv f_i(\mathbf{h}^{m-1}, \mathbf{x}^{-m}) \quad \forall i \leq m-1.$$

Total differentiation yields

$$\frac{\partial h_i^{m-1}}{\partial x_m} = \sum_{j=1}^{m-1} \frac{\partial f_i}{\partial h_j} \cdot \frac{\partial h_j^{m-1}}{\partial x_m} + \frac{\partial f_i}{\partial x_{m+1}}.$$

In matrix notation this gives

$$\begin{aligned} \frac{\partial \mathbf{h}^{m-1}}{\partial x_m} &= \mathbf{J}_m \frac{\partial \mathbf{h}^{m-1}}{\partial x_m} - \mathbf{A}_{Cm} \\ \Rightarrow \frac{\partial \mathbf{h}^{m-1}}{\partial x_m} &= -\mathbf{A}_{m-1}^{-1} \mathbf{A}_{Cm} \end{aligned}$$

so Equation (11) becomes

$$\frac{dg_m}{dx_m} = \mathbf{A}_{Rm} \mathbf{A}_{m-1}^{-1} \mathbf{A}_{Cm}.$$

Finally, note that

$$\mathbf{A}_{Rm} \mathbf{A}_{m-1}^{-1} \mathbf{A}_{Cm} = 1 - \frac{|\mathbf{A}_m|}{|\mathbf{A}_{m-1}|},$$

which gives us Equation (9).³

Ω

The conditions for an iterative quasi-contraction mapping can now be stated in terms of the determinants of the \mathbf{A}_m matrices:

Theorem 6:

a) If there is an ordering of players, indexed by 1..n, such that for some $e \in (0,1)$

$$|\mathbf{A}^m| \geq e > 0 \quad \forall m \in \{2..n\} \tag{12}$$

then the best-response function exhibits an iterative quasi-contraction mapping.

b) If, in addition, we have

$$\frac{|\mathbf{A}^m|}{|\mathbf{A}^{m-1}|} \leq 2 - e < 2 \quad \forall m \in \{2..n\}. \tag{13}$$

³ This is a special case of the general result for partitioned matrices that

$$\begin{vmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{B} & \mathbf{D} \end{vmatrix} = |\mathbf{A}| |\mathbf{D} - \mathbf{B}\mathbf{A}^{-1}\mathbf{C}|$$

for square matrices \mathbf{A} and \mathbf{D} , where \mathbf{A} is non-singular. For a proof of this result, see, for example, Rao (1965, p28).

then the best-response function exhibits an iterative quasi-contraction mapping.

Proof:

Follows automatically from Theorems 3 and 4.

Ω

Condition (12) gives the general uniqueness condition of this paper—that the game has a unique equilibrium if the matrix $I-J$ is positive definite. In the remainder of the paper, we show that this condition encompasses and extends existing uniqueness conditions.

6. Relationship to Other Uniqueness Conditions.

A. Rosen's Theorem:

The best-known paper providing a generic set of sufficient conditions for games of the form analysed here is Rosen (1965). Rosen considers a very general game structure in which the strategy space for any player can be conditional on the strategy chosen by another (as could happen in a coalition game). In the special case, however, where the player's strategy spaces are orthogonal to each other, i.e. the class of games considered in this paper, Rosen's sufficient condition is that as follows:

If there exists a diagonal matrix R , with diagonal terms $r_{ii} > 0 \forall i$ such that the symmetric matrix $(RA) + (RA)'$ is positive definite, then there is a unique equilibrium.

The main result of this paper is more general than this in two ways. First, when $n > 2$, it is possible to construct examples where the conditions for Theorem 6 holds but not Rosen's theorem. (When $n = 2$, the matrix conditions are equivalent.) Second, Rosen first establishes existence by means of the Kakutani fixed-point theorem, and thus requires each player's strategy space be bounded, closed and convex, none of which are required by the version of the contraction mapping theorem used here.

B. Cournot Games.

There are many papers giving conditions for uniqueness in a Cournot quantity-setting game. These include Szidarovszky and Yakowitz (1977), Gaudet and Salant (1991), and Long and Soubeyran (2000). All three of these papers provide conditions which imply that the best-response functions of firms are negatively sloped, along with other conditions required to bound the set of prices over which demand is positive. As shown by the following, theorem, by using the general uniqueness theorem in this paper, one only needs to require non-positively-sloped best-response functions; the bounding conditions are not necessary.

Theorem 7:

Let A_n be a square matrix with $a_{ii} = 1 \forall i$ and

$$a_{ij} = a_i \in (-1, 0] \quad \forall j \neq i, \quad \forall i.$$

Then $|A_n| > 0$.

Proof:

Given in the Appendix .

Ω

C. Row-Sum Conditions:

Cachon and Netessine (2004) show that a sufficient condition for the aggregate best-response function, \mathbf{f} , to exhibit a contraction mapping is that

$$\sum_{j \neq i} \left| \frac{\partial f_i}{\partial x_j} \right| < 1 \quad \text{or} \quad \sum_{i \neq j} \left| \frac{\partial f_j}{\partial x_i} \right| < 1.$$

That is, \mathbf{f} exhibits a contraction mapping if the sum of the absolute values of the off-diagonal elements in the Jacobean matrix be less than one in each row or in each column. This result is established by showing that a function has a contraction mapping if the largest eigenvalue of the Jacobean matrix is less than one, and then using a result of Horn and Johnson (1996), that this will hold if the maximum row sum or the maximum column sum is less than one. Although this approach is very different from ours, it is easy to show that this condition meets our requirement for there to be an iterative quasi-contraction mapping:

Theorem 8:

Let A_n be a square matrix with $a_{ii} = 1 \forall i$ and $\sum_{j \neq i} |a_{ij}| < 1$. Then $|A_m| > 0 \forall m \leq n$.

Proof:

Given in the Appendix.

Ω

References:

- Cachon, G. and S. Netessine (2004), "Game Theory in Supply Chain Analysis", in *Supply Chain Analysis in the E-Business Era*, D. Simchi-Levi, S.D. Wu and M. Shen, Eds., Kluwer.
- Gaudet, G. and S. W. Salant (1991), "Uniqueness of cournot equilibrium: New results from Old Methods", *Review of Economic Studies*, 58(2): 399–404.
- N. V. Long and A. Soubeyran (2000), "Existence and Uniqueness of Cournot Equilibrium: A Contraction Mapping Approach", *Economic Letters*. 2000, Vol.67: 345-8.
- Rao, C.R. (1965), *Linear Statistical Inference and its Applications*. John Wiley and Sons: New York.
- Rosen, J.B. (1965), "Existence and uniqueness of equilibrium points for concave N-person games", *Econometrica*, Vol.33: 520-33.
- Szidarovszky, F. and Yakowitz, S. (1977), "New Proof of the Existence and Uniqueness of the Cournot Equilibrium". *International Economic Review*, 18:787-89.
- Takayama, A. (1985), *Mathematical Economics* 2nd edition, Cambridge University Press: Cambridge.

Appendix

A. Proof of Theorem 7.

Let Ω_n be the set of $n \times n$ matrices, A_n , satisfying the following properties:

- a) $a_{ii} = 1 \quad \forall i$
- b) $\exists \bar{a} \in (0,1)$ s.t. $a_{ij} = a_i \in [0, \bar{a}] \quad \forall i, i \neq j$,

We want to show that

$$|A_n| > 0 \quad \forall A_n \in \Omega_n.$$

Proof:

The proof is by induction. The proposition is clearly true for $n=1$ and $n=2$. Now assume that there is some $\bar{n} > 2$ such that the proposition holds for all $n < \bar{n}$. we will show that it then holds for $n = \bar{n}$.

First note that if $a_i = 0$ for any i , then $|A_n| = |A_n^{-(i,i)}|$, and, since $A_n \in \Omega_n \Rightarrow A_n^{-(i,i)} \in \Omega_{n-1}$, the result holds by the induction assumption. We shall therefore only consider the case where $a_i > 0 \quad \forall i$.

Now define the matrix $B_m(A_n, b_m)$, which is derived from some $n \times n$ matrix, A_n , by replacing a_{mm} with b_m . The determinant of this matrix is

$$|B_m(A_n, b_m)| = |A_n| - (a_{mm} - b_m) |A_n^{-(m,m)}|.$$

We can show the following result:

Lemma 1:

If $|A_m| > 0$ for all $A_m \in \Omega_m$, for all $m \leq n$, then $|B_m(A_m, a_m)| \geq 0 \quad \forall A_m \in \Omega_m$.

That is, changing the diagonal term in the bottom row from unity to be the same as the other $m-1$ terms in that row will not change the sign of the determinant of the matrix from positive to negative.

Proof:

The proof is by contradiction. Assume that $|B_m(A_m, a_m)| < 0$. Note that

$$|B(A_m, b_m)| = |A_m| - (1 - b_m) |A_{m-1}|.$$

Since $|A_{m-1}|$ is positive by assumption, $|B(A_m, 1, b_m)|$ is continuous and monotone increasing in b_m . Further, since $|B(A_m, 1, 1)| > 0$, by the intermediate value theorem there must exist some $\hat{b}_m \in (a_m, 1)$ such that $|B(A_m, 1, \hat{b}_m)| = 0$. Now multiply the m th row of $B(A_m, 1, \hat{b}_m)$ by $1/\hat{b}_m$. Since \hat{b}_m is positive, this leaves the sign of the determinant unchanged. But it produces a matrix that has exactly the form we are considering and so must have a positive determinant. This establishes the contradiction.

Now consider the matrix $B_{n-1}(B_n(A_n, \mathbf{a}_n), \mathbf{a}_{n-1})$ —that is, the matrix found by replacing the diagonal terms in each of the last two rows with the entry for the off-diagonal terms in those rows. Since the n th row is then a multiple of the $(n-1)$ th row,

$$|B_{n-1}(B_n(A_n, \mathbf{a}_n), \mathbf{a}_{n-1})| = 0.$$

Using Equation (?) above, we can also write

$$\begin{aligned} |B_{n-1}(B_n(A_n, \mathbf{a}_n), \mathbf{a}_{n-1})| &= |B_n(A_n, \mathbf{a}_n)| - (1 - a_{n-1}) |B_n(A_n, \mathbf{a}_n)^{-(m-1, m-1)}| \\ &= |A_n| - (1 - a_n) |A_n^{-(n, n)}| - (1 - a_{n-1}) |B_n(A_n, \mathbf{a}_n)^{-(m-1, m-1)}|. \end{aligned}$$

By the induction base, we have

$$|A_n^{-(n, n)}| > 0 \quad \forall A_n \in \Omega_n,$$

and by the induction base and Lemma 1, we have

$$|B_n(A_n, \mathbf{a}_n)^{-(m-1, m-1)}| \geq 0 \quad \forall A_n \in \Omega_n.$$

We therefore have

$$|A_n| > |B_{n-1}(B_n(A_n, \mathbf{a}_n), \mathbf{a}_{n-1})| = 0,$$

which establishes the result.